

Expanded BRFSS, 2003

Bronx Locality

Table of Contents:

Optional Module 1: Adult Asthma History..... 1
Optional Module 5: Diabetes..... 2
Optional Module 7: Disability/Quality of Life..... 6

Optional Module 10: Health Care Coverage..... 10
Optional Module 13: Injury Control - Falls..... 12
Optional Module 19: Tobacco Consumption..... 13
Optional Module 20: Tobacco - Media..... 14
Optional Module 21: Tobacco - other products..... 22
Optional Module 22: Tobacco - Work site ETS..... 23
Optional Module 23: Seat Belts..... 25
Optional Module 24: Weight Control..... 27

New York State Department of Health

Expanded BRFSS, 2003 - New York State Department of Health

Have had an Asthma Episode or Attack in the Past 12 Months, among Those with Asthma

Bronx

Demographic Groups	n ¹	Yes % ²	n	No %	C.I. ³
Total	36	55.0	33	45.0	14.0

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

Check Blood for Glucose or Sugar at Least Once per Day among Those with Diabetes

Bronx

Demographic Groups	n ¹	Yes % ²	n	No %	C.I. ³
Total	26	35.3	40	64.7	13.6

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

Check Feet for Sores or Irritations at Least Once per Day among Those with Diabetes

Bronx

Demographic Groups	n ¹	Yes % ²	n	No %	C.I. ³
Total	42	65.2	21	34.8	14.2

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

Had Feet Checked by a Medical Professional in Past Twelve Months, among Those with Diabetes

Bronx

Demographic Groups	n ¹	Yes % ²	n	No %	C.I. ³
Total	47	77.6	15	22.4	12.1

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

Had Dilated Eye Exam in Past 12 Months, among Those with Diabetes

Bronx

Demographic Groups	n ¹	Yes % ²	n	No %	C.I. ³
Total	49	77.5	14	22.5	12.5

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

Adults with a Disability

Bronx

Demographic Groups	n¹	Yes %²	n	No %	C.I.³
Total	118	17.3	494	82.7	3.4
Gender					
Male	29	14.1	158	85.9	5.4
Female	89	20.0	336	80.0	4.2
Age					
18-34	19	9.0	161	91.0	4.5
35-54	52	18.7	214	81.3	5.3
≥55	47	28.6	110	71.4	8.2
Education					
≤High School	73	20.7	249	79.3	5.1
>High School	44	13.4	240	86.6	4.3

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

Adults with a Disability Requiring Help

Bronx

Demographic Groups	Disability need help			Disability no help needed			No Disability		
	n ¹	% ²	C.I. ³	n	%	C.I.	n	%	C.I.
Total	61	9.5	2.6	57	7.8	2.3	494	82.7	3.4
Gender									
Male	12	6.4	3.9	17	7.7	4.0	158	85.9	5.4
Female	49	12.1	3.5	40	7.9	2.7	336	80.0	4.2
Age									
18-64	50	8.7	2.6	42	6.4	2.4	439	84.8	3.4
≥65	11	15.3	9.5	15	16.9	8.6	46	67.8	11.8
Education									
≤High School	43	12.6	4.1	30	8.0	3.4	249	79.3	5.1
>High School	17	5.6	2.9	27	7.8	3.3	240	86.6	4.3

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

This page left intentionally blank

This page left intentionally blank

Expanded BRFSS, 2003 - New York State Department of Health

Main Reason for Not Having Health Care Coverage

Bronx

Demographic Groups	n ¹	Changed or lost job % ²	C.I. ³	n	Could not afford %	C.I.	n	Other %	C.I.
Total	12	8.7	5.9	27	31.6	11.4	66	59.6	11.7

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

No Health Care Coverage for Past Two Years Among Uninsured

Bronx

Demographic Groups	n ¹	Yes % ²	n	No %	C.I. ³
Total	43	44.0	60	56.0	11.9

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

Had a Fall in Past Three Months

Bronx

Demographic Groups	n ¹	No fall % ²	C.I. ³	n	Fall with injury %	C.I.	n	Fall no injury %	C.I.
Total	237	84.3	5.0	18	7.1	3.6	25	8.6	3.8

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

Number of Cigarettes Smoked per day among Everyday Smokers

Bronx

Demographic Groups	n¹	1-10 cigarettes %²	n	≥11 cigarettes %	C.I.³
Total	51	63.0	32	37.0	12.2

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

Heard, Read or Seen Information about the Effects of Tobacco from Television

Bronx

Demographic Groups	n¹	Yes %²	n	No %	C.I.³
Total	416	68.8	177	31.2	4.6
Gender					
Male	118	63.8	63	36.2	8.2
Female	298	72.8	114	27.2	5.0
Age					
18-34	136	75.8	39	24.2	8.0
35-54	182	66.2	76	33.8	7.0
≥55	93	63.1	59	36.9	9.2
Education					
≤High School	215	68.0	92	32.0	6.5
>High School	198	70.8	82	29.2	6.4

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

Heard, Read or Seen Information about the Effects of Tobacco from Radio

Bronx

Demographic Groups	n¹	Yes %²	n	No %	C.I.³
Total	230	39.2	356	60.8	4.8
Gender					
Male	72	39.6	108	60.4	8.3
Female	158	38.9	248	61.1	5.5
Age					
18-34	74	40.5	99	59.5	8.9
35-54	102	40.8	151	59.2	7.2
≥55	49	34.2	103	65.8	9.1
Education					
≤High School	128	42.4	180	57.6	6.8
>High School	101	36.2	171	63.8	6.7

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

Heard, Read or Seen Information about the Effects of Tobacco from Billboards

Bronx

Demographic Groups	n¹	Yes %²	n	No %	C.I.³
Total	278	50.7	312	49.3	4.9
Gender					
Male	86	52.0	94	48.0	8.5
Female	192	49.6	218	50.4	5.6
Age					
18-34	109	66.8	67	33.2	8.3
35-54	117	46.8	136	53.2	7.3
≥55	49	33.0	104	67.0	9.0
Education					
≤High School	147	51.5	160	48.5	6.9
>High School	130	50.5	147	49.5	7.0

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

Heard, Read or Seen Information about the Effects of Tobacco from Buses

Bronx

Demographic Groups	n ¹	Yes % ²	n	No %	C.I. ³
Total	218	36.4	371	63.6	4.7

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

Heard, Read or Seen Information about the Effects of Tobacco from Magazines

Bronx

Demographic Groups	n¹	Yes %²	n	No %	C.I.³
Total	317	51.9	270	48.1	4.9
Gender					
Male	93	50.6	86	49.4	8.5
Female	224	53.0	184	47.0	5.6
Age					
18-34	101	57.2	73	42.8	9.1
35-54	144	53.8	112	46.2	7.2
≥55	69	41.3	81	58.7	9.4
Education					
≤High School	160	49.8	148	50.2	6.9
>High School	156	55.5	117	44.5	7.0

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

Heard, Read or Seen Information about the Effects of Tobacco from a Website

Bronx

Demographic Groups	n ¹	Yes % ²	n	No %	C.I. ³
Total	80	15.1	517	84.9	3.5

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

Heard, Read or Seen Information about the Effects of Tobacco from Movie Theater

Bronx

Demographic Groups	n ¹	Yes % ²	n	No %	C.I. ³
Total	117	22.6	473	77.4	4.3

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

Heard or Seen Anti-Tobacco Advertisement that Refers to the NYS Quitline in Past 30 Days

Bronx

Demographic Groups	n ¹	Yes % ²	n	No %	C.I. ³
Total	301	53.9	275	46.1	5.0
Gender					
Male	100	55.7	79	44.3	8.5
Female	201	52.4	196	47.6	5.7
Age					
18-34	109	62.7	64	37.3	8.9
35-54	127	49.7	124	50.3	7.3
≥55	63	48.2	83	51.8	9.8
Education					
≤High School	160	56.3	141	43.7	6.8
>High School	139	52.3	130	47.7	7.1

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

Ever Used or Tried Any Smokeless Tobacco Products such as Chewing Tobacco or Snuff

Bronx

Demographic Groups	n ¹	Yes % ²	n	No %	C.I. ³
Total	27	4.4	568	95.6	2.1

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

Work Indoors Most of the Time

Bronx

Demographic Groups	n¹	Yes %²	n	No %	C.I.³
Total	259	77.6	66	22.4	5.6

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

Smoking Policy at Place of Work among Indoor Workers**Bronx**

Demographic Groups	n¹	Not allowed anywhere %²	C.I.³	n	Allowed in some or all places %	C.I.	n	No official policy %	C.I.
Total	210	75.5	7.5	30	17.1	7.0	17	7.4	4.2
Gender									
Male	57	64.1	13.1	15	24.9	12.6	11	10.9	7.9
Female	153	86.1	6.2	15	9.8	5.5	6	4.1	3.2
Age									
18-34	61	69.2	14.1	12	22.0	13.3	5	8.8	8.5
35-54	113	79.3	8.5	15	14.2	8.0	10	6.5	4.1
≥55	32	86.2	12.6	2	7.5	10.0	2	6.3	8.5
Education									
≤High School	76	70.1	13.1	13	23.0	12.8	6	6.9	6.3
>High School	134	81.6	7.7	17	12.9	7.2	10	5.5	3.5

¹Percents based on row denominators of less than 50 are unstable and should be used with caution²Weighted Percent³95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

Use Seatbelt when Ride or Drive in a Car (Always, Nearly Always or Never Ride in a Car)

Bronx

Demographic Groups	n ¹	Yes % ²	n	No %	C.I. ³
Total	518	86.7	73	13.3	3.4

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

Use Seatbelt when Ride or Drive in a Car (Always or Never ride in a Car)

Bronx

Demographic Groups	n¹	Yes %²	n	No %	C.I.³
Total	458	74.9	133	25.1	4.3
Gender					
Male	127	70.3	52	29.7	7.6
Female	331	78.5	81	21.5	4.8
Age					
18-64	393	73.5	122	26.5	4.8
≥65	58	82.1	11	17.9	10.5
Education					
≤High School	230	74.0	78	26.0	5.9
>High School	226	77.7	51	22.3	6.1

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

Trying to Lose Weight

Bronx

Demographic Groups	n¹	Yes %²	n	No %	C.I.³
Total	291	45.7	300	54.3	4.9
Gender					
Male	72	38.4	106	61.6	8.2
Female	219	51.5	194	48.5	5.6
Age					
18-34	76	37.2	99	62.8	8.4
35-54	136	49.9	121	50.1	7.2
≥55	76	52.0	76	48.0	9.5
Education					
≤High School	152	45.5	157	54.5	6.8
>High School	135	45.7	141	54.3	7.0

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

Trying to Maintain Current Weight of Those not Trying to Lose Weight

Bronx

Demographic Groups	n¹	Yes %²	n	No %	C.I.³
Total	187	62.9	113	37.1	6.7
Gender					
Male	63	58.8	44	41.2	11.0
Female	124	67.1	69	32.9	7.4
Age					
18-34	57	58.4	41	41.6	12.0
35-54	82	69.1	40	30.9	9.6
≥55	47	62.4	29	37.6	12.8
Education					
≤High School	96	65.3	61	34.7	9.1
>High School	90	61.3	51	38.7	9.7

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

Eating Fewer Calories or Less Fat to Lose or Maintain Weight

Bronx

Demographic Groups	n¹	Yes %²	n	No %	C.I.³
Total	370	77.3	102	22.7	4.7
Gender					
Male	100	75.9	32	24.1	8.6
Female	270	78.3	70	21.7	5.2
Age					
18-34	99	73.4	33	26.6	9.4
35-54	166	77.4	49	22.6	6.4
≥55	101	81.8	20	18.2	8.9
Education					
≤High School	191	76.5	53	23.5	6.7
>High School	175	78.5	48	21.5	6.3

¹Percents based on row denominators of less than 50 are unstable and should be used with caution

²Weighted Percent

³95% Confidence Interval

Expanded BRFSS, 2003 - New York State Department of Health

Using Physical Exercise to Lose or Maintain Weight***Bronx***

Demographic Groups	n¹	Yes %²	n	No %	C.I.³
Total	295	60.7	182	39.3	5.4
Gender					
Male	88	64.6	47	35.4	9.5
Female	207	58.0	135	42.0	6.2
Age					
18-34	94	69.3	39	30.7	9.8
35-54	136	62.2	81	37.8	7.6
≥55	62	46.3	61	53.7	10.5
Education					
≤High School	130	53.0	118	47.0	7.6
>High School	162	70.8	62	29.2	7.0

¹Percents based on row denominators of less than 50 are unstable and should be used with caution²Weighted Percent³95% Confidence Interval